

Abundances and Correlations of Structures Beyond Galaxy Clusters

Wen Xu¹, Li-Zhi Fang², Zugan Deng³

ABSTRACT

We investigated the structures on scales beyond the typical clusters of galaxies. These structures are crucial to understand the cosmic gravitational clustering in the pre-virialized stage, or quasilinear regime. Based on the multi-resolution analysis of the discrete wavelet transformation, we got statistical available ensembles of r_{cl} -clusters, i.e. the structures on scale r_{cl} , in the range $1 \leq r_{cl} \leq 24 h^{-1}$ Mpc for both N-body simulation and the IRAS 1.2 Jy galaxy survey samples. If the mass-to-light ratio on scales larger than clusters asymptotically reaches a constant, we found that the abundances and correlations of these IRAS r_{cl} -clusters to be basically consistent with the predictions of the flat low-density CDM model (LCDM) and the open CDM model (OCDM), except the model-predicted abundance of $r_{cl}=24 h^{-1}$ Mpc clusters seems to be higher than IRAS data. The standard CDM (SCDM) gives too much power, and too weak correlations on all the scales. For a given r_{cl} , the amplitude of two-point correlation function of r_{cl} -clusters is increasing with their richness. However, for a given richness (defined by the mean separation of neighbor objects), the clustering strengths of both simulation and observation sample are found to be declining with r_{cl} when r_{cl} is larger than 3 - 4 h^{-1} Mpc. Therefore, the “universal” increase of the correlation amplitude with the scale of objects from galaxies, groups, to poor, rich clusters is broken down for structures of $r_{cl} > 3 - 4 h^{-1}$ Mpc. Supercluster should not be a member of the “universal” increase family.

Subject headings: cosmology: theory – galaxies: clusters: general – large-scale structure of universe

1. Introduction

The study of clusters of galaxies is among the most fruitful topics in cosmology. It provides rich information on large scale structures, significant for understanding the nature of formation and

¹ Beijing Astronomical Observatory, Beijing 100080, China; also currently at Department of Physics and Astronomy, Arizona State University, Tempe, AZ 85287

²Department of Physics, University of Arizona, Tucson, AZ 85721

³Department of Physics, Graduate School, CAS, P.O. Box 3908, Beijing, China

evolution of cosmic structures on scales of $\sim 1 - 10$ Mpc. In particular, cosmological parameters such as the density parameter Ω_0 , the mass fluctuation amplitude σ_8 and even the cosmological constant Ω_λ can be effectively constrained by observed properties of clusters. For instance, low mass density cold dark matter (CDM) models, $\Omega_0 \simeq 0 - 0.3$, are found to be generally favored by the abundance and two-point correlation function of clusters (Jing & Fang 1994; Eke et al. 1996; Viana & Liddle 1996; Carlberg et al. 1997; Bahcall, Fan & Cen 1997; Croft et al. 1997).

However, the redshift and peculiar velocity data on scales larger than clusters, such as IRAS galaxy survey, yield $\Omega_0 \simeq 0.3 - 1.5$, with a trend towards $\Omega_0 \simeq 1$ if moderate biasing is taken into account (e.g. Dekel 1994; Strauss & Willick 1995 for summary of the results). Moreover, structures on scales larger than typical clusters are pre-virialized, or depend on the quasilinear evolution of cosmic clustering. Namely, the larger-than-cluster (LTC) objects are in a quantitatively different dynamical evolution stage from the objects on scales equal to or less than Abell rich clusters which are virialized, or nearly virialized. Thus, one cannot estimate the clustering behavior of larger-than-cluster (LTC) objects by the results of clusters. Therefore, it is worth to investigate whether the low mass density CDM cosmogony is still favored by the abundance and two-point correlation function of structures on scales larger than clusters. This is the goal of this paper.

A curious thing is that we know of some observed LTC structures, like superclusters, and we know the dynamics of structure formation of popular models, but at the moment have not put them together. This is largely because the lack of a well-defined ensemble of LTCs. Structures on scales larger than clusters generally are not “regular” in both configuration and velocity spaces. There is no well-defined dynamical criterion to classify and identify these structures, and therefore, statistical approaches have not yet been carefully pursued.

This problem can be solved by a *complete* decomposition. Since a complete decomposition does not lose information, any “irregular” objects or distributions can be precisely represented by the coefficients of a complete decomposition. Thus, the model-observation confrontation can be done with a decomposed representation if the decomposition can reasonably work for both observed and model-predicted distributions. Recently it has been shown that a statistical analysis can indeed be performed for un-virialized large scale structures by a complete multi-resolution analysis of the discrete wavelet transformation (*DWT*). It has successfully identified structures on scales from 1 to about $20 h^{-1}$ Mpc, and revealed the scaling behavior of LTC clustering (Xu, Fang & Wu 1998, hereafter Paper I).

This paper is organized as follows. In §2 we present the abundances of IRAS r_{cl} -clusters and compare with the model predictions. §3 tests the models with the IRAS two-point correlation functions of the LTC structures. Finally, conclusion is in §4.

2. Abundances of r_{cl} -clusters

2.1. Mass functions of r_{cl} -clusters from simulation samples

The model samples of mass distributions used in this paper are generated by N-body simulations with the P³M code developed by Y.P. Jing (Jing & Fang 1994, Jing et al. 1995). Three popular models of the CDM cosmogony, i.e. the standard CDM (SCDM), the flat low-density CDM (LCDM) models and the open CDM model (OCDM), are employed. The Hubble constant h ($\equiv H_0/(100 \text{ km/sec/Mpc})$), mass density Ω_0 , cosmological constant Ω_λ and the normalization of the power spectra of the density perturbation σ_8 are chosen to be (0.5, 1.0, 0.0, 0.62) for SCDM, (0.75, 0.3, 0.7, 1.0) for LCDM, and (0.75, 0.3, 0.0, 1.0) for OCDM. The simulation parameters are: 1.) box size $L = 310 h^{-1} \text{ Mpc}$; 2.) number of simulation particles $N_p = 64^3$ and 3.) the effective force resolution $\eta = 0.24 h^{-1} \text{ Mpc}$. In this case, a particle has mass of $3.14 \times 10^{13} \Omega_M h^{-1} M_\odot$, which is small enough to resolve reliably objects with mass $M \geq 5.5 \times 10^{14} h^{-1} M_\odot$. We produced 8 realizations for each model. The mass of a r_{cl} -cluster is measured within a sphere of radius r_{cl} .

The method of finding clusters on scales r_{cl} by discrete wavelet transformation (DWT) has been developed in Paper I (about the technical aspects of the DWT multi-scale identification, see also Fang & Thews 1998). We identified clusters on scales $r_{cl} = 0.75, 1.5, 3, 6, 12$, and $24 h^{-1} \text{ Mpc}$ from the simulated samples. A result is shown in Fig. 1. It plots the top 25 massive r_{cl} -clusters for each scale of $r_{cl} = 0.75, 1.5, 3, 6, 12$, and $24 h^{-1} \text{ Mpc}$, from one realization of OCDM simulation with volume $310^3 h^{-3} \text{ Mpc}^3$.

Observationally, supercluster is defined as cluster of rich clusters (top r_{ab} -clusters). In Fig. 1, some r_{ab} -clusters are located together to form large scale clumps or filaments. These structures are obviously superclusters. Fig. 1 also shows, however, that some of the top 25 $r_{cl} = 24 h^{-1} \text{ Mpc}$ halos contain nothing of $r_{cl} \leq 24 h^{-1} \text{ Mpc}$ massive halos. This is, although the mass and length scales of these $r_{cl} = 24 h^{-1} \text{ Mpc}$ clusters are the same as superclusters, they do not contain any of the top 25 clusters on smaller scales. This result indicates that the identification of superclusters by “cluster of rich clusters” may miss some very massive objects on scale $r_{cl} = 24 h^{-1} \text{ Mpc}$. Recently, it has been found that the dynamical properties of superclusters are not only determined by rich clusters, but also by numerous galaxies in the intra-cluster regions (Small et al. 1998). This also implies that one should directly identify LTC structures from matter or galaxy distributions, instead of using only rich clusters.

With the multi-scale identification of simulation samples, we have model-predicted mass function of r_{cl} -clusters, $n(> M, r_{cl})$, which is the number density of r_{cl} -clusters with mass larger than M . The mass functions of clusters with $r_{cl} = 1.5, 3, 6$ and $12 h^{-1} \text{ Mpc}$ in models of the SCDM, LCDM and OCDM are shown in Fig. 2. The mass function of clusters on Abell scale, i.e. $r_{cl} = r_{ab} = 1.5 h^{-1} \text{ Mpc}$, is the same as that given by *friends-of-friends* identification (Jing & Fang 1994). We have also calculated mass functions of $r_{cl} = 24 h^{-1} \text{ Mpc}$ clusters, which are not shown in Fig. 2, but will be employed below. The statistical errors of the mass functions of r_{cl} -cluster are mainly caused by Poisson errors. The middle part of the mass function is usually more reliable, as all the Poisson errors are relatively small in this range. In Fig. 2, this middle

range corresponds roughly to $10^{-4} < n(> M, r_{cl}) < 10^{-7} h^3 \text{ Mpc}^{-3}$.

Since each cluster now is characterized by two parameters M and r_{cl} , it is inconvenient to define richness by mass alone. Instead, the relative richness can be described by the mean neighbor separation d between the clusters considered. The larger the d , the higher the richness.

2.2. r_{cl} -clusters from IRAS samples

We have also applied *DWT* multi-resolution analysis on real samples of the redshift survey of the Infrared Astronomical Satellite (IRAS) galaxies with flux limit of $f_{60} \geq 1.2 \text{ Jy}$ (Fisher et al. 1995). The IRAS 1.2 Jy sample contains 5313 IRAS galaxies, covering the whole sky with depth of about $150h^{-1} \text{ Mpc}$.

The sampling rate of the IRAS sample is low. Therefore, the IRAS clusters identified on Abell scale r_{ab} contain much fewer galaxies than optical clusters. This gives rise to higher Poisson errors. However, larger r_{cl} clusters contain more members, and Poisson error is lower. The IRAS sample can give statistically significant ensemble of $r_{cl} > r_{ab}$ clusters. Moreover, the IRAS sample consists of less early type of galaxies, and more late type. Namely, it has less galaxies in the center of r_{ab} -clusters, but more field galaxies. This is rather a good point for studying structures beyond clusters, as late type galaxies probably are more fair tracers of mass field on larger scales.

To minimize the radial effects, we've divided the data into redshift intervals, or the velocity intervals of [2500, 5000], [5000, 7500], [7500, 10000] and [10000, 12500] km s^{-1} . We then identify r_{cl} -clusters in each interval. The details of cluster-finding algorithm has been discussed in Paper I.

It has been shown that IRAS galaxies basically trace the local large scale structures. Some individual clusters and superclusters from optical surveys have also been identified from the IRAS data (Fisher et al. 1995; Webster et al. 1997). We checked this point further by cross-identification between the Abell clusters and the multi-scale identified r_{cl} clusters of the IRAS galaxies. Using the position data (l , b , and z) collected in the compilation of the Abell catalog (Abell, Corwin & Olowin 1989), one can find 18 rich clusters (with galaxy counts > 30) in the range [5000, 7500] km s^{-1} . They are A194, A347, A400, A426, A539, A569, A779, A1367, A1656, A2870, A2877, A2911, A2933, A3389, A3537, A3656, A3706, A3879.

Since the maximum positioning uncertainty of our cluster finding algorithm is about $3 h^{-1} \text{ Mpc}$ in 2-d, and 2000 km s^{-1} in radial, we choose the matching radius r between the projections of centers of the cross-identified clusters to be less than $3 h^{-1} \text{ Mpc}$, and the radial separation d of the centers less than 2000 km s^{-1} . In this case, 7 out of the 18 Abell clusters are cross-identified with IRAS $r_{cl} = 3h^{-1} \text{ Mpc}$ clusters. These are A347, A426, A539, A1367, A3389, A3656, A3706. The overall confidence level of this cross-identification can be estimated by a binomial distribution function $B(k, n, p)$ that a random distribution would generate $\geq k$ cross-identifications from n realizations when the chance alignment probability is known as p . When overlapping is not

important, $p = N\pi r^2 d/V$, where N and V are the total number of clusters and the volume of the redshift shell, respectively. For the above-mentioned Abell-IRAS cluster cross-identification the overall confidence level is 3σ ($\sim 99.7\%$).

The difference between 7 and 18 is due partially to the low sampling rate of the IRAS, i.e. the higher Poisson errors. It is not a surprise that the richness of individual clusters estimated from IRAS may have large difference from optical estimate. This cross-identification seems to show that the Abell clusters may be underestimated by the IRAS identification by a factor about 2.6. However, we also did the cross-identification in the opposite direction, i.e. to identify the 16 top IRAS clusters in the shell $[5000, 7500]$ km s $^{-1}$ with Abell clusters. The result is: 5 out of the 16 IRAS clusters have their Abell counterparts. Namely, the IRAS clusters may be underestimated by the Abell clusters by a factor about 3. Therefore, statistically, both Abell and IRAS clusters trace the same density field. If we adjust the matching radius r and separation d , we would find that the number of Abell-IRAS cross-identified clusters increases from the 7 of 18 (3σ confidence level), to 13 (2σ) and 15 (1σ). This shows that the Abell and IRAS clusters provide about the same description of the massive halos on scale r_{ab} .

Using the 7 better cross-identified $r_{cl} = 3h^{-1}$ Mpc IRAS-Abell clusters, one can calculate the mean ratio of N_{IRAS}/N_{abell} , where $N_{IRAS} = N_g(1.5)/\phi(z)$, N_{abell} is the number of galaxies of the considered Abell cluster, $N_g(1.5)$ is the observed number of IRAS galaxies within radius $1.5 h^{-1}$ Mpc around the cluster center, and $\phi(z)$ the IRAS selection function (Fisher et al. 1995). We found that the median value $N_{IRAS}/N_{abell} = 7.03$. On the other hand, the mass-to-number ratio of Abell clusters on average is $M_{abell}/N_{abell} = 0.6 \times 10^{13} h^{-1} M_\odot$, where M_{abell} is the mass of the Abell clusters (Bahcall and Cen 1993). Thus, the total mass of the IRAS clusters M can be represented as

$$M = AN_{IRAS}, \quad (1)$$

where the number-mass conversion coefficient A is

$$A = 10^{11.93} M_\odot. \quad (2)$$

2.3. Mass function of IRAS r_{ab} -clusters

Considering the luminosity is proportional to the number of galaxies, the number-mass conversion relation eq.(1) essentially is the mass-to-light ratio for IRAS r_{ab} -clusters. Moreover, the mass-to-light ratio for Abell clusters is independent of the total mass of the clusters (Bahcall & Cen 1993). Therefore, it seems to be reasonable to employ eq.(1) to find the total mass of IRAS r_{ab} -clusters.

The total number of member galaxies, N_{IRAS} , is given by $N_{IRAS} = N_g/\phi(z)$, where $\phi(z)$ is the selection function (Fisher et al. 1995). Thus, we have

$$M = AN_g/\phi(z). \quad (3)$$

Using eq.(3), the mass function of the IRAS r_{ab} -clusters can be found from their abundance $n_{nc}(> N_g, r_{cl})$ by

$$n_{IRAS}(> M, r_{ab}) = n_{nc}(> N_g = M\phi(z)/A, r_{ab}). \quad (4)$$

Using the $n_{nc}(> N_g, r_{ab})$ given by the IRAS galaxies in the shell [5000, 7500] km s⁻¹, we calculated the mass function of the IRAS r_{ab} -clusters $n_{IRAS}(> M, r_{ab})$. We found that $n_{IRAS}(> M, r_{ab})$ is in very good agreement with the mass function of optical and X-ray clusters within Abell radius, $n_{op}(> M, r_{ab})$ (Bahcall and Cen 1992), especially for rich clusters of $M > 10^{14} h^{-1} M_{\odot}$.

We have also calculated the mass function $n_{IRAS}(> M, r_{ab})$ using the IRAS data of the shells [2500, 5000], [7500, 10000] and [10000, 12500] km s⁻¹. The resulted $n_{IRAS}(> M, r_{ab})$ is also basically in agreement with $n_{op}(> M, r_{ab})$. The goodness of this agreement can be described by the number-mass conversion coefficient A . In order to have a best fitting of $n_{IRAS}(> M, r_{ab})$ with $n_{op}(> M, r_{ab})$, we find $\log A = 12.07 \pm 0.21$ for shell [2500, 5000] km s⁻¹, 11.93 ± 0.15 for [5000, 7500], 11.71 ± 0.32 for [7500, 10000], 11.63 ± 0.41 for [10000, 12500]. Here the errors include both the Poisson errors of n_{IRAS} and n_{op} . A shows a slight decrease from lower velocity shell to higher velocity. This might be caused either by the uncertainty of the selection function, or by a very weak redshift variation of IRAS galaxies with respect to optical clusters.

The conversion coefficients A from all the redshift shells are the same as the A of eq.(2) within errors, and their differences are not larger than a factor of 3. This is, in terms of mass function of r_{ab} -clusters, the matter distributions traced by the IRAS galaxies is about the same as optical and X-ray clusters on large scales. This result seems to be a little surprising. It is known that the luminous IRAS galaxies with heavy dust lane do not appear at the densest regions, namely the clustering of infrared galaxies is weaker than that of optical galaxies at the core region of rich clusters. Therefore, the matter distributions represented by IRAS galaxies should be less clustering than that traced by optical galaxies on small scales. In other words, the mass-to-light ratio should actually be higher for IRAS clusters at the core region. However, the mass function given by eq.(4) matches well with that of the optical clusters. This is because the mass function eq.(4) measures statistically the mean number of collapsed massive r_{ab} -halos over a large volume. It is actually a measure of clustering features on scales equal to or larger than clusters. It thus less depends on the physical process within the core region. In a word, the IRAS galaxies may cause underestimates of individual masses of some richest clusters on smaller scales, while for large scales, IRAS galaxies give about the same statistically features of mass field as optical objects.

2.4. Model comparisons of abundances of IRAS r_{cl} -clusters

Similar to eq.(4), one may derive the mass function of IRAS r_{cl} -clusters, $n_{IRAS}(> M, r_{cl})$, from their number-count function $n_{nc}(> N_g, r_{cl})$ by

$$n_{IRAS}(> M, r_{cl}) = n_{nc}(> N_g = M\phi(z)/A, r_{cl}). \quad (5)$$

The problem is yet how to determine A at scale $r_{cl} > r_{ab}$.

From eq.(1), A essentially is the mass-to-light ratio of the system on scale r_{ab} . The derived value of A for r_{ab} -clusters corresponds to a median mass-to-light ratio of $M/L \simeq 300 \pm 100 h(M/L)_{\odot}$. As mentioned above, r_{ab} -clusters are dominated by elliptical galaxies, while the larger structures have higher fraction of spiral galaxies. The mass-to-light ratio or A on scales $r_{cl} > r_{ab}$ may be smaller than r_{ab} . The observed Virgo cluster infall motion, however, indicates that M/L ratio doesn't show a decrease with scale, while appears to flatten and remain approximately constant on scales larger than $0.2 h^{-1}$ Mpc (Bahcall, Lubin & Dorman 1995). This result is strengthened by a direct measure of M/L of the Corona Borealis supercluster (Small et al. 1998). It gives $M/L \simeq 560 h(M/L)_{\odot}$. Namely, the M/L ratio on scale $\sim 20 h^{-1}$ Mpc is not lower, but even higher than that of rich clusters by a factor about 1.9. This implies that the M/L ratio is weakly scale-dependent from scales of 1 to about $20 h^{-1}$ Mpc. The weak dependence of M/L on scale is also supported by the following fact: the segregation between elliptical and spiral galaxies of APM Bright Galaxies Catalogue (BGC), as well as the ratio between the second order bias factors of these APM-BGC elliptical and spiral galaxies, is found to be basically scale-independent from 1 to $20 h^{-1}$ Mpc (Loveday et al. 1995, Fang, Deng & Xia, 1998). It would be reasonable to assume that the elliptical-spiral mix doesn't cause a serious scale-dependence of the mass-to-light ratio on large scales, and A is scale-independent within a factor of 3.

Using the A for each shell (§2.3), we calculated the mass functions of IRAS clusters with $r_{cl} = 1.5, 3, 6, 12$ and $24 h^{-1}$ Mpc for data of the velocity shells $[2500, 5000]$, $[5000, 7500]$, $[7500, 10000]$ and $[10000, 12500]$ km s^{-1} . The errors of the IRAS cluster mass functions are estimated by the same way as that of optical clusters. The vertical error bars in Figs. 3 - 6 are the Poisson errors of counting the clusters in the limited volume of a shell. The horizontal error bars are the Poisson errors of mass estimation of each clusters. This Poisson error is generally smaller for larger r_{cl} , as more galaxies are contained in larger r_{cl} clusters. Therefore, at larger r_{cl} the IRAS data can set effective constraints.

Since the ratio between mass functions at r_{cl} and r_{ab} is independent of the value A , Figs. 3 - 6 essentially are tests of the r_{cl} -dependence of mass functions of r_{cl} clusters. Despite the errors of these figures are large, Figs. 3 and 4 have shown that all $n(> M, r_{cl})$ in the range of $r_{cl} \leq 12 h^{-1}$ Mpc and in the two shells $[2500, 5000]$ and $[5000, 7500]$ km s^{-1} are in good agreement with the predictions of the models of LCDM and OCDM. The results of $24 h^{-1}$ Mpc in these two shells (which have not directly been shown in Figs 3 and 4) are also basically consistent with LCDM and OCDM. The dynamical status of r_{ab} -clusters is very different from $r_{cl} \gg r_{ab}$ clusters. The former are most likely virialized, the later are not. Therefore, Figs. 3 and 4 indicate that the models of OCDM and LCDM are still in good shape for clustering in pre-virialization regimes.

For shells of $[7500, 10000]$ and $[10000, 12500]$ km s^{-1} , Figs. 5 and 6 show that the LCDM and OCDM can fit with the data of $r_{cl} \leq 6 h^{-1}$ Mpc. Yet, the IRAS mass functions on scales of $r_{cl} = 12$ and $24 h^{-1}$ Mpc are lower than the predictions of LCDM and OCDM. The deficit of

IRAS $r_{cl} = 24 h^{-1}$ Mpc clusters may result from the poorness of the current data at > 7500 km s^{-1} . Since the overdensity of the large scale objects is relatively low, their detection will be poorer than small scales. Moreover, not all predicted $r_{cl} = 12, 24 h^{-1}$ Mpc clusters contain top rich r_{ab} -clusters (Fig.1), and therefore, some of these structures will be overlooked if the completeness of samples is not high enough.

Figs. 3-6 clearly show a systematic and significant lower of the IRAS mass functions in all velocity shells than the predictions of SCDM. SCDM gives too many massive clusters on *all* scales of $r_{cl} \leq 24 h^{-1}$ Mpc. It is interesting to note that if all IRAS mass functions are shifted along the $\log M$ axis by a factor of about 0.60, the SCDM model will be able to survive. The shifting factor 0.60 requires A to be larger than our derived value by a factor $10^{0.6} \simeq 4$, and the corresponding ratio M/L should be as high as $4 \times 300 \simeq 1200h(M/L)_{\odot}$. Thus, the SCDM can not be saved if the uncertainty of A is less than a factor of 4.

3. Two Point Correlation Functions of r_{cl} -Clusters

3.1. d - dependence of correlations

It has been known for a long time that the two-point cluster correlation function $\xi_{cc}(r)$ follows the same power law as galaxies, i.e. $\xi_{cc}(r) = (r/r_0)^{-\gamma}$ and $\gamma \sim 1.8$, but has much larger correlation length r_0 than galaxies. Moreover, the correlation length r_0 defined by $\xi(r_0) \equiv 1$ is found to be systematically increasing with the cluster richness. An empirically “universal” scaling $r_0 = 0.4d$, d being the mean neighbor separation as employed to describe the cluster richness (§2.1), was proposed to fit with the correlations of galaxies, QSOs, group of galaxies, X-ray and Abell clusters and even superclusters (e.g. Bahcall & Soneira 1983; Bahcall & West 1992 and references therein). Because the sizes of these objects, from galaxies to superclusters, are systematically increasing by at least two orders, it is unclear whether various objects with very different scales can really be fitted with one “universal” scaling.

This problem can be solved by studying the two-point correlations of clusters with different r_{cl} . Using samples of r_{cl} -clusters from N-body simulation, we found that the two-point correlation functions of r_{cl} -clusters can be well fitted with a power law $\xi(r) = (r/r_0)^{-\gamma}$ for all considered r_{cl} and d . The index γ is in the range of 1.8 - 2.0 for the LCDM and OCDM, and 1.9-2.1 for the SCDM. No indication shows that the index γ would significantly vary with either r_{cl} or d .

The r_0 - d relations for r_{cl} -clusters are calculated. The results of the OCDM are plotted in Fig. 7. This figure shows that the r_0 - d lines for $r_{cl} = 0.75, 1.5$ and $3 h^{-1}$ Mpc almost coincide. This is, in the range of $r_{cl} \leq 3 h^{-1}$ Mpc, the r_0 - d relation is “universal”, independent of the size of structures. The model SCDM and LCDM show similar “universal” r_0 - d relations.

However, the r_0 - d relation is substantially dependent on r_{cl} if $r_{cl} > 3 h^{-1}$ Mpc. Despite the correlation length r_0 is still increasing with richness d , the lines $r_0 - d$ are lower for larger

r_{cl} . Therefore, superclusters are not a member of the universal r_0 - d relation of $r_{cl} \leq 3 h^{-1}$ Mpc clusters.

For clusters of r_{cl} larger than $12 h^{-1}$ Mpc, their r_0 is comparable to and even less than r_{cl} . In this case, r_0 is actually determined by an extrapolation of the two-point correlation function $\xi(r)$ from range $r > r_{cl}$ to $r < r_{cl}$, and solving $\xi(r_0) = 1$. This method might be susceptible to large error. Nevertheless, it is reliable that lines of $r_0 - d$ are significantly lower than the $r_0 - d$ of smaller r_{cl} clusters.

In redshift space the correlation length will increase by no more than $2h^{-1}$ Mpc, and the index γ becomes slightly larger. Our overall conclusions on correlation functions will not change due to these slight differences.

3.2. r_{cl} -dependence of correlations

Fig.8 plots the correlation length r_0 as a function of r_{cl} for a given d . This figure, actually, is an alternative representation of Fig.7. It shows that for a given d the correlation length r_0 is almost invariant for r_{cl} equal to or less than $3 h^{-1}$ Mpc, but slowly decreasing with r_{cl} when r_{cl} is larger than about $4 h^{-1}$ Mpc. This is a common feature for all the three models.

Theoretically, the decrease of r_0 with r_{cl} is expected. Fig. 1 shows that most halos of $r_{cl} = 0.75, 1.5$ or $3 h^{-1}$ Mpc coincide to form a 0.75-1.5-3 clouds-in-clouds. This is because most halos of $r_{cl} \leq 2 h^{-1}$ Mpc are virialized. Therefore, the detection of r_{cl} -clusters with $r_{cl} \leq 3 h^{-1}$ Mpc will find almost the same objects. In other words, the ensembles of $r_{cl} = 0.75, 1.5$ or $3 h^{-1}$ Mpc clusters actually consist of the same objects, and therefore, they have the same correlations, and the universal r_0 - d relations. However, on scales larger than $4 h^{-1}$ Mpc, more and more halos are less developed, and their correlations are lower. Therefore, the “universal” increase of the correlation amplitudes with scale from galaxies, groups to clusters should be broken down for $r_{cl} > 3 - 4 h^{-1}$ Mpc clusters. Superclusters are not members of the “universal” scaling family.

3.3. Correlation lengths of IRAS clusters

With the samples of the IRAS r_{cl} -clusters, we calculated the correlation length r_0 by fitting the observed data with $\xi(r) = (r/r_0)^{-1.8}$. The correlation lengths r_0 are measured from data of different velocity shells. The results of r_0 vs. d are plotted in Fig. 9. The data points shown in Fig. 9 are binned in d . The errors are given by the variance of the correlation strengths at each bin of d . For $r_{cl} = r_{ab}$, the values of r_0 found from the IRAS samples are in agreement with those from APM sample (Bahcall & Cen 1992; Croft et al. 1997). Fig. 9 shows, obviously, that the r_0 - d relation from IRAS samples of $r_{cl} = 1.5$ to $12 h^{-1}$ Mpc favors the models of OCDM and LCDM. The SCDM-predicted r_0 - d curves are systematically lower than the IRAS results for all considered

d.

We also tested the models by the r_{cl} -dependence of r_0 . Fig.10 gives the r_0 - r_{cl} relations from the simulation samples and IRAS data. The errors of the IRAS data are also large. But the observed data show once again that the LCDM and OCDM are basically consistent with the IRAS data, while the SCDM-predicted r_0 is too low to match with the observation. More importantly, it is clear to see the slow decrease of r_0 with the increase of r_{cl} in all models. The r_0 decrease is an indicator of the transform from quasilinear to non-linear régimes.

4. Conclusions and discussions

Structures on scales equal to and larger than typical clusters have been identified by the *DWT* decomposition. This method is shown to be able to systematically and uniformly identify structures on various scales, especially those of larger than the size of typical clusters. Taking this advantage, one can produce statistically available ensembles of r_{cl} -clusters from simulation samples and observational data.

To describe the dynamics of the cosmic clustering on scales beyond clusters, we introduced several new statistical measures: the bivariant abundance function $n(> M, r_{cl})$, the r_{cl} -dependence of abundance, the r_{cl} -dependence of correlations, etc. These statistics are useful to reveal the features of pre-virialized regime of cosmic gravitational clustering, and to discriminate among models of structure formation, especially on large scales.

The identification of LTCs is effective to extract information from sparsely distributed data like the IRAS galaxies. The IRAS 1.2 Jy galaxies have low sampling rate compared to optical galaxies and are not found at cluster centers. They are not very effective to be employed for identification of traditional r_{ab} -clusters. However, the IRAS clusters with $r_{cl} > 3 h^{-1}$ Mpc can already set effective constraints on models, if the mass-to-light ratio for structures traced by IRAS galaxies is assumed to asymptotically reach a constant on scales larger than Abell radius. Therefore, IRAS galaxies can provide the same information of large scale structures as optical and X-ray galaxies, if the considered scales are larger than the scale of morphological segregation of IRAS galaxies that no dusty spirals like IRAS sources exist at the immediate neighbourhood of densest regions.

The multi-resolution analysis of the IRAS sample concludes that the abundances $n(> M, r_{cl})$ and correlations of objects on scales $r_{cl} = 1$ to $24 h^{-1}$ Mpc can be described by LCDM and OCDM, while the SCDM is in difficulty. Namely, the abundances and two-point correlation functions of structures on scales as large as about $20 h^{-1}$ Mpc favor a low-density universe if M/L ratio is assumed to be flat over these scales.

We would like to thank an anonymous referee for a detailed report which improves the

presentation of the paper. Wen Xu thanks the World Laboratory for a scholarship. This project was done when WX was visiting at Physics Department, University of Arizona.

REFERENCES

- Abell, G. O., Corwin, Jr. H. G. & Olowin, R. P. 1989, ApJS, 70, 1.
- Bahcall, N. A. & Cen, R. Y. 1993, ApJ, 407, L49.
- Bahcall, N. A., Fan, X. & Cen, R. Y. 1997, ApJ, 485, L53.
- Bahcall, N. A., Lubin, L. M. & Dorman, V. 1995, ApJ, 447, L81.
- Bahcall, N. A., & Soneira, R. M. 1983, ApJ, 270, 20.
- Carlberg, R. G., Morris, S. M., Yee, H. K. C. & Ellingson, E. 1997, ApJ, 479, L19.
- Croft, R. A. C., Dalton, G. B., Efstathiou, G., Sutherland, W. J. & Maddox, S. J. 1997, MNRAS, 291, 305.
- Dekel, A., 1994, ARA&A, 32, 371.
- Eke, V. R., Cole, S., Frenk, C. S. & Navarro, J. F. 1996, MNRAS, 281, 703.
- Fang, L. Z., Deng, Z. G. & Xia, X. Y. 1998, ApJ, 506, 53.
- Fang, L. Z. & Thews R. 1998, *Wavelts in Physics*, World Scientific, Singapore.
- Fisher, K. B. et al. 1995, ApJS, 100, 69.
- Jing, Y. P. & Fang, L. Z. 1994, ApJ, 432, 438.
- Jing, Y. P., Mo, H. J., Börner, G. & Fang, L. Z. 1995, MNRAS, 274, 417.
- Loveday, L., Maddox, S. J., Efstathiou, G. & Peterson, B. A. 1995, ApJ, 442, 457.
- Small, T., Ma, C.-P., Sargent, W., Hamilton, D. 1998, ApJ, 492, 45
- Strauss, M.A. & Willick, J.A. 1995, Phys. Rep., 261, 271
- Viana, P. P. & Liddle, A. R., 1996, MNRAS, 281, 323.
- Webster, M., Lahav, O., & Fisher, K. 1997, MNRAS, 287, 425.
- Xu, W., Fang, L. Z. & Wu, X. P. 1998, ApJ, 508, 472. (Paper I)

Fig. 1.— A projected distribution of r_{cl} -clusters identified from a realization of OCDM N-body simulation within box $310 h^{-3} \text{ Mpc}^3$. It shows the top 25 of the richest clusters for each r_{cl} . The legends of the symbols are: plus for 0.75, diamond for 1.5, circle for 3, triangle for 6, square for

12 and giant circle for $24 h^{-1}$ Mpc. The linear sizes of symbols are roughly equal to their r_{cl} , but slightly amplified for small r_{cl} for clarity.

Fig. 2.— Mass functions of clusters identified with radii $r_{cl} = 1.5, 3.0, 6.0$ and $12 h^{-1}$ Mpc for SCDM, LCDM and OCDM models at redshift $z = 0$. $n(> M, r_{cl})$ is the number density of r_{cl} -clusters with mass larger than M . M is in unit of $h^{-1} M_{\odot}$.

Fig. 3.— Mass functions of IRAS r_{cl} -clusters. n is in unit of $h^3 \text{ Mpc}^{-3}$. The horizontal errors are Poisson errors from the number counting of galaxies in a given cluster, and the vertical errors are the Poisson errors from counting the clusters. The model predictions of SCDM, LCDM, OCDM are shown as dashed lines, thin solid lines, and thick solid lines, respectively. Observed data are from the IRAS sample in the velocity shell $[2500, 5000] \text{ km s}^{-1}$.

Fig. 4.— The same as Fig. 3, but the data are from shell $[5000, 7500] \text{ km s}^{-1}$.

Fig. 5.— The same as Fig. 3, but the data are from shell $[7500, 10000] \text{ km s}^{-1}$.

Fig. 6.— The same as Fig. 3, but the data are from shell $[10000, 12500] \text{ km s}^{-1}$.

Fig. 7.— The r_0 - d relations for clusters with different r_{cl} in the OCDM model. The error bars of the curves of $r_{cl} = 6$ and $12 h^{-1}$ Mpc are from the scattering among realizations of simulation.

Fig. 8.— The r_0 - r_{cl} relations at different richness for models of OCDM, LCDM, SCDM.

Fig. 9.— The r_0 - d relations for $r_{cl} = 1.5, 3, 6, 12 h^{-1}$ Mpc. The dashed, thin solid, and thick solid lines are, respectively, the simulation results of models SCDM, LCDM and OCDM. The observed data are from the averages over the velocity shells of the IRAS galaxies.

Fig. 10.— The r_0 - r_{cl} relations for $d = 25, 50, 70 h^{-1}$ Mpc. The dashed, thin solid, and thick solid lines are, respectively, the simulation results of models SCDM, LCDM and OCDM. The observed data are from the averages over the velocity shells of the IRAS galaxies.



















